

A Combinatorial Classification of Skew Schur Functions

Peter McNamara
Bucknell University

Joint work with Stephanie van Willigenburg

Special Session on Algebraic Combinatorics
AMS Sectional Meeting, Fayetteville, AR
3 November 2006

Slides and paper available from
www.facstaff.bucknell.edu/pm040/

When are Two Skew Schur Functions Equal?

Peter McNamara
Bucknell University

Joint work with Stephanie van Willigenburg

Special Session on Algebraic Combinatorics
AMS Sectional Meeting, Fayetteville, AR
3 November 2006

Slides and paper available from
www.facstaff.bucknell.edu/pm040/

- ▶ Background: skew Schur functions
- ▶ Recent work on skew Schur function equality
- ▶ Skew Schur equivalence
- ▶ Composition of skew diagrams, main results
- ▶ Conjectures, open problems


Schur functions

▶ Partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$

▶ Young diagram.

Example:

$$\lambda = (4, 4, 3, 1)$$



Schur functions


▶ Partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$

▶ Young diagram.

Example:

$$\lambda = (4, 4, 3, 1)$$

▶ Semistandard Young tableau (SSYT)



The Schur function s_λ in the variables $x = (x_1, x_2, \dots)$ is then defined by

$$s_\lambda = \sum_{\text{SSYT } T} x_1^{\#1\text{'s in } T} x_2^{\#2\text{'s in } T} \dots$$

Example

$$s_{4431} = x_1 x_3^2 x_4^4 x_5 x_6^2 x_7 x_9 + \dots$$

Skew Schur functions

▶ Partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_\ell)$


▶ μ fits inside λ .

▶ Young diagram.

Example:

$$\lambda/\mu = (4, 4, 3, 1)/(3, 1)$$

▶ Semistandard Young tableau (SSYT)



The **skew** Schur function $s_{\lambda/\mu}$ in the variables $x = (x_1, x_2, \dots)$ is then defined by

$$s_{\lambda/\mu} = \sum_{\text{SSYT } T} x_1^{\#1\text{'s in } T} x_2^{\#2\text{'s in } T} \dots$$

Example

$$s_{4431/31} = x_4^3 x_5 x_6^2 x_7 x_9 + \dots$$

- ▶ Skew Schur functions are symmetric in the variables $x = (x_1, x_2, \dots)$.
- ▶ The Schur functions form a basis for the algebra of symmetric functions (over \mathbb{Q} , say).
- ▶ Connections with Algebraic Geometry, Representation Theory


Big Question: When is $s_{\lambda/\alpha} = s_{\mu/\beta}$?

Big Question: When is $s_{\lambda/\alpha} = s_{\mu/\beta}$?

- ▶ Lou Billera, Hugh Thomas, Steph van Willigenburg (2004):


Big Question: When is $s_{\lambda/\alpha} = s_{\mu/\beta}$?

- ▶ Lou Billera, Hugh Thomas, Steph van Willigenburg (2004):




Big Question: When is $s_{\lambda/\alpha} = s_{\mu/\beta}$?

- ▶ Lou Billera, Hugh Thomas, Steph van Willigenburg (2004):



Complete classification of equality of ribbon Schur functions



- ▶ HDL II: Vic Reiner, Kristin Shaw, Steph van Willigenburg (2006):
 - ▶ The more general setting of binomial syzygies

$$cS_{D_1} S_{D_2} \cdots S_{D_m} = c' S_{D'_1} S_{D'_2} \cdots S_{D'_n}$$

is equivalent to understanding equalities among connected skew diagrams.


- ▶ 3 operations for generating skew diagrams with equal skew Schur functions.
- ▶ Necessary conditions, but of a different flavor.

- ▶ HDL III: McN., Steph van Willigenburg (2006):
 - ▶ An operation that encompasses the three operations of HDL II.
 - ▶ Theorem that generalizes all previous results.
Explains the 6 missing equivalences from HDL II.
 - ▶ Conjecture for necessary and sufficient conditions for $s_{\lambda/\alpha} = s_{\mu/\beta}$.
Reflects classification of HDL I for ribbons.

Skew diagrams (skew shapes) D , E .

If $s_D = s_E$, we will write $D \sim E$.

Example



We want to classify all equivalence classes, thereby classifying all skew Schur functions.

The basic building block

EC2, Exercise 7.56(a) [2-]

Theorem

$D \sim D^*$, where D^* denotes D rotated by 180° .

The basic building block

EC2, Exercise 7.56(a) [2-]

Theorem

$D \sim D^*$, where D^* denotes D rotated by 180° .

Goal: Use this equivalence to build other skew equivalences.

The basic building block

EC2, Exercise 7.56(a) [2-]

Theorem

$D \sim D^*$, where D^* denotes D rotated by 180° .

Goal: Use this equivalence to build other skew equivalences.

Where we're headed:


Theorem

Suppose we have skew diagrams D , D' and E satisfying certain assumptions. If $D \sim D'$ then


$$D' \circ_W E \sim D \circ_W E \sim D \circ_{W^*} E^*.$$

Main definition: composition of skew diagrams.


Composition of skew diagrams




Composition of skew diagrams




Composition of skew diagrams



Composition of skew diagrams




Composition of skew diagrams



Theorem [McN., van Willigenburg] *If $D \sim D'$, then*


$$D' \circ E \sim D \circ E \sim D \circ E^*.$$



Amalgamated Compositions

Actually, the previous slide was just a warm-up....


A skew diagram W *lies in the top* of a skew diagram E if W appears as a connected subdiagram of E that includes the northeasternmost cell of E .



Amalgamated Compositions

Actually, the previous slide was just a warm-up....

A skew diagram W *lies in the top* of a skew diagram E if W appears as a connected subdiagram of E that includes the northeasternmost cell of E .




Similarly, W *lies in the bottom* of E .

Our interest: W lies in both the top and bottom of E . We write $E = WOW$.

Amalgamated Compositions

Actually, the previous slide was just a warm-up....

A skew diagram W *lies in the top* of a skew diagram E if W appears as a connected subdiagram of E that includes the northeasternmost cell of E .




Similarly, W *lies in the bottom* of E .

Our interest: W lies in both the top and bottom of E . We write $E = WOW$.


Hypotheses: (inspired by hypotheses of RSvW)

1. W is maximal given its set of diagonals.
2. W_{ne} and W_{sw} are separated by at least one diagonal.
3. $E \setminus W_{ne}$ and $E \setminus W_{sw}$ are both connected skew diagrams.


Amalgamated Compositions




Amalgamated Compositions



Amalgamated Compositions




Amalgamated Compositions




15 boxes: first of the non-RSvW examples

Amalgamated Compositions




15 boxes: first of the non-RSvW examples

If $W = \emptyset$, we get the regular compositions:




What are the results?

Construction of \overline{W} and \overline{O} :




What are the results?

Construction of \overline{W} and \overline{O} :



What are the results?


Construction of \overline{W} and \overline{O} :



Hypothesis 4. \overline{W} is never adjacent to \overline{O} .

What are the results?

Construction of \overline{W} and \overline{O} :




Hypothesis 4. \overline{W} is never adjacent to \overline{O} .

Conjecture. Suppose we have skew diagrams D, D' with $D \sim D'$ and $E = WOW$ satisfying Hypotheses 1-4, then

$$D' \circ_W E \sim D \circ_W E \sim D \circ_{W^*} E^*.$$

What are the results?

Construction of \overline{W} and \overline{O} :




Hypothesis 4. \overline{W} is never adjacent to \overline{O} .

Conjecture. Suppose we have skew diagrams D, D' with $D \sim D'$ and $E = WOW$ satisfying Hypotheses 1-4, then

$$D' \circ_W E \sim D \circ_W E \sim D \circ_{W^*} E^*.$$

Hypothesis 5. In $E = WOW$, at least one copy of W has just one cell adjacent to O .



What are the results?


Theorem.[McN., van Willigenburg] Suppose we have skew diagrams D, D' with $D \sim D'$ and $E = WOW$ satisfying Hypotheses 1-5, then

$$D' \circ_W E \sim D \circ_W E \sim D \circ_{W^*} E^*.$$

What are the results?

Theorem.[McN., van Willigenburg] Suppose we have skew diagrams D, D' with $D \sim D'$ and $E = \text{WOW}$ satisfying Hypotheses 1-5, then

$$D' \circ_W E \sim D \circ_W E \sim D \circ_{W^*} E^*.$$



15 boxes: second of the non-RSvW examples

The hard part: An expression for $s_{D \circ_W E}$ in terms of s_D , s_E , $s_{\overline{W}}$, $s_{\overline{O}}$:


$$s_{D \circ_W E} (s_{\overline{W}})^{|\widehat{D}|} (s_{\overline{O}})^{|\widetilde{D}|} = \pm (s_D \circ_W s_E).$$

The easy part: The blue portion is invariant if we replace D by D' when $D' \sim D$. Similarly, can replace E by E^* .


Proof of expression uses:

- ▶ Hamel-Goulden determinants. See paper of Chen, Yan, Yang.
- ▶ Sylvester's Determinantal Identity.

- ▶ Removing Hypothesis 5.



$D \circ_W E$ has 23 boxes, and $D \circ_W E \sim D^* \circ_W E$:



Main open problem

Theorem. [McN, van Willigenburg]

Skew diagrams E_1, E_2, \dots, E_r

$E_i = W_i O_i W_i$ satisfies Hypotheses 1-5

E'_i and W'_i denote either E_i and W_i , or E_i^* and W_i^* .

Then

$$((\cdots (E_1 \circ_{W_2} E_2) \circ_{W_3} E_3) \cdots) \circ_{W_r} E_r \sim ((\cdots (E'_1 \circ_{W'_2} E'_2) \circ_{W'_3} E'_3) \cdots) \circ_{W_r} E'_r.$$

Main open problem

Theorem. [McN, van Willigenburg]

Skew diagrams E_1, E_2, \dots, E_r

$E_i = W_i O_i W_i$ satisfies Hypotheses 1-5

E'_i and W'_i denote either E_i and W_i , or E_i^* and W_i^* .

Then

$$((\cdots (E_1 \circ_{W_2} E_2) \circ_{W_3} E_3) \cdots) \circ_{W_r} E_r \sim ((\cdots (E'_1 \circ_{W'_2} E'_2) \circ_{W'_3} E'_3) \cdots) \circ_{W_r} E'_r.$$

Conjecture. [McN, van Willigenburg; inspired by main result of BTvW]

Two skew diagrams E and E' satisfy $E \sim E'$ if and only if, for some r ,

$$\begin{aligned} E &= ((\cdots (E_1 \circ_{W_2} E_2) \circ_{W_3} E_3) \cdots) \circ_{W_r} E_r \\ E' &= ((\cdots (E'_1 \circ_{W'_2} E'_2) \circ_{W'_3} E'_3) \cdots) \circ_{W_r} E'_r, \text{ where} \end{aligned}$$

- $E_i = W_i O_i W_i$ satisfies Hypotheses 1-4 for all i ,
- E'_i and W'_i denote either E_i and W_i , or E_i^* and W_i^* .

Main open problem

Theorem. [McN, van Willigenburg]

Skew diagrams E_1, E_2, \dots, E_r

$E_i = W_i O_i W_i$ satisfies Hypotheses 1-5

E'_i and W'_i denote either E_i and W_i , or E_i^* and W_i^* .

Then

$$((\cdots (E_1 \circ_{W_2} E_2) \circ_{W_3} E_3) \cdots) \circ_{W_r} E_r \sim ((\cdots (E'_1 \circ_{W'_2} E'_2) \circ_{W'_3} E'_3) \cdots) \circ_{W_r} E'_r.$$

Conjecture. [McN, van Willigenburg; inspired by main result of BTvW]

Two skew diagrams E and E' satisfy $E \sim E'$ if and only if, for some r ,

$$\begin{aligned} E &= ((\cdots (E_1 \circ_{W_2} E_2) \circ_{W_3} E_3) \cdots) \circ_{W_r} E_r \\ E' &= ((\cdots (E'_1 \circ_{W'_2} E'_2) \circ_{W'_3} E'_3) \cdots) \circ_{W_r} E'_r, \text{ where} \end{aligned}$$

- $E_i = W_i O_i W_i$ satisfies Hypotheses 1-4 for all i ,
- E'_i and W'_i denote either E_i and W_i , or E_i^* and W_i^* .

True for $n \leq 19$.

- ▶ A definition of skew diagram composition. Encompasses the composition, amalgamated composition and staircase operations of RSvW.
- ▶ Theorem that generalizes all previous results. In particular, explains the 6 missing equivalences from HDL II.
- ▶ Conjecture for necessary and sufficient conditions for $E \sim E'$.